



Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Texas


Monday, 04 January 2016

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Monday, December 28, 2015



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from December 26 to January 1: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at:

<http://www.tpwd.state.tx.us/landwater/water/envconcerns/hab/redtide/status.phtml>

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive:

<http://tidesandcurrents.noaa.gov/hab/bulletins.html>

Conditions Report

There is currently no indication of *Karenia brevis* (commonly known as Texas red tide) along the coast of Texas. No respiratory irritation is expected Monday, January 4 through Monday, January 11.

Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.


Analysis

Sampling from the Texas A&M University's Imaging FlowCytobot, located on the Port Aransas ship channel, indicates that *Karenia brevis* is 'not present' (TAMU; 1/1-4). For information on area shellfish restrictions, contact the Texas Department of State Health Services.

Recent MODIS Aqua imagery (1/3, shown left) is partially obscured by clouds along-and offshore the Texas coast from the Galveston Island region to south of the Rio Grande, limiting analysis. Patches of elevated to high chlorophyll (2-16 $\mu\text{g/L}$) are visible along-and offshore the Texas coast from Sabine Pass to Galveston Island. Elevated chlorophyll is not indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast.

Forecast models based on predicted near-surface currents indicate a potential maximum transport of 60 km south from the Port Aransas region from January 3-7.


Kavanaugh, Lalime



Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

Wind Analysis

Port Aransas to Baffin Bay: Northeast winds (10-15kn, 5-8m/s) today becoming east winds after midnight through Tuesday night. Southeast winds (5-10kn, 3-5m/s) Wednesday becoming southwest winds after midnight. Northwest to west winds (5-10kn) Thursday becoming south winds after midnight. Southwest winds (5kn, 3m/s) Friday becoming southeast winds (5-10kn) Friday night.



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).